



Tendon transfers for irreparable rotator cuff tears: An update

Jeremie M. Axe

Shoulder dysfunction in the setting of irreparable rotator cuff tears (RCTs) can be treated successfully with different types of tendon transfer:

- Latissimus dorsi transfer for irreparable posterosuperior RCTs works best for young, active patients with an intact subscapularis, no pseudoparalysis or previous surgery, and a functioning teres minor.
- A more anatomical transfer for irreparable posterosuperior RCTs is a lower trapezius transfer, and early results are promising.
- Isolated irreparable tears of the subscapularis can be successfully managed with pectoralis major tendon transfer with a concentric humeral head. However, restricted external rotation (ER) may occur, depending on technique.
- Pectoralis minor transfer can successfully address combination irreparable tears of the upper border subscapularis and the supraspinatus without significant loss of ER.
- Rotator cuff arthropathy with ER lag benefits most from a reverse total shoulder arthroplasty and a combination latissimus dorsi and teres major transfer (LDTMT) regardless of patient age.

Keywords: tendon transfers; shoulder; irreparable; latissimus dorsi; rotator cuff arthropathy

Cite this article: Axe JM. Tendon transfers for irreparable rotator cuff tears: An update. *EFORT Open Rev* 2016;1:18-24.

Tendon transfers of the shoulder can be technically challenging, but if performed well can significantly improve the quality of a patient's life. Surgical skill is essential, but just as important is proper patient selection. Despite advances in surgical techniques and equipment, failure to heal may occur after rotator cuff tear (RCT) repair and may not be amenable to revision surgery. This review, structured in three sections, is designed to investigate current literature on transfers, over the last five years, on irreparable posterosuperior RCTs, anterosuperior RCTs and transfers with reverse total shoulder arthroplasty (RTSA).

The following sections describe several different transfers in detail; however, a brief anatomical refresher

regarding innervations of the muscles involved is in order. Essentially, four muscles comprise the rotator cuff: the supraspinatus, the infraspinatus, the teres minor and the subscapularis. The supraspinatus and infraspinatus muscles are innervated by the suprascapular nerve. The teres minor muscle is innervated by the axillary nerve and the subscapularis muscle by the upper and lower subscapular nerves. The latissimus dorsi muscle is innervated by the thoracodorsal nerve and is the largest muscle in the back, and functions as an adductor, extender and internal rotator of the arm. The trapezius muscle has three sets of fibres and is innervated by the spinal accessory nerve or 11th cranial nerve. The teres major muscle is innervated by the lower subscapular nerve and, like the latissimus, is an adductor and internal rotator of the arm. The medial pectoral nerve innervates both the pectoralis major and minor, and the lateral pectoral nerve innervates just the pectoralis major.

Irreparable posterosuperior RCTs

Reports of massive posterosuperior rotator cuff repair failures range from 21% to 91%^{1,2,3} and revision failure rates are significantly higher.⁴ Recurrent tears typically occur within the first six months following primary fixation.^{5,6} Symptoms of re-tear at two years include: impaired overhead function, increased pain, limited passive movement, loss of strength, and lower overall satisfaction with shoulder function.⁴ Complications also arise when performing revision rotator cuff repairs.⁷ Fortunately, when a posterosuperior RCT tear is deemed irreparable, there are several options for the patient.

Gerber originally defined an irreparable posterosuperior RCT as the inability to achieve fixation in $\leq 60^\circ$ of abduction despite adequate releases.⁸ This also includes at least stage 3 fatty infiltration of the supraspinatus and infraspinatus, an acromiohumeral distance of < 7 mm on imaging⁸ or static migration of the humeral head.⁹ Excluding shoulder arthroplasty, options considered for irreparable recurrent tears are debridement with possible partial repair, augmentation with scaffolding, or tendon transfers.¹⁰ There are several possible major tendon transfers for irreparable posterosuperior RCTs. These include the latissimus dorsi transfer (LDT) with or without the teres major (LDTMT) and the lower trapezius transfer (LTT).

LDT

The earliest and most studied transfer is the LDT, originally described by Gerber in 1988.⁸ It has the potential of a 33 cm excursion when fully released from surrounding attachments.¹¹ The intention in performing a LDT is to restore humeral head depression and external rotation (ER) in the young, active population without glenohumeral (GH) arthritis or significant static migration of the humeral head. It essentially reconstitutes the posterior force couple.⁹ As with most surgeries, there is appropriate patient selection with respect to demographics, the physical exam., and radiographic imaging.

Approximately 70% of LDTs are performed on men. However, there is no objective outcome difference regarding gender.¹² Namdari et al¹² suggested that it is difficult to assess whether hand dominance and occupation are clinically significant, as corresponding reports were underpowered and inconsistent. While no age cap has been described, a recent systematic review noted a mean age of 59 years for LDT.¹² LDTs in the presence of prior rotator cuff repair have resulted in poorer patient outcomes,^{13,14,15,16,17} especially with reference to post-operative constant scores.

Physical examination of the patient is critical. The subscapularis tendon should be intact and functioning, as forward elevation drastically decreases with insufficiency.^{18,19,20} Additionally, GH stability increases with an intact subscapularis in the setting of an LDT.²¹ Positive drop-arm²² and ER lag tests²³ are typical findings. Supple shoulder movement is essential – specifically, passive forward flexion (FF) and abduction $\geq 80^\circ$.²⁴ A pseudoparalytic shoulder has been demonstrated to correlate with poor outcomes.²⁵ Axillary nerve lesions and deltoid insufficiencies are contra-indications.

Imaging plays a role in the decision-making process. Standard shoulder radiographs and MRI are common modalities. The patient cannot have radiographic indications of GH arthritis, and limited, Hamada stage 1 or 2, rotator cuff arthropathy.¹⁹ This implies no acetabulisation of the acromion.²⁶ The irreparable supraspinatus and infraspinatus tendons are typically torn with retraction to the level of the glenoid (Patte stage 3),²⁷ with fatty infiltration Goutallier grade 3^{28,29} and/or significant atrophy.³⁰ Subscapularis tears with grade 3 or higher Goutallier atrophy,^{28,29} and/or $\geq 50\%$ tear of the upper border³¹ should also be excluded.¹⁹ Atrophy of the teres minor assessed pre-operatively by MRI was performed by several authors.^{32,16,17} It was concluded that fatty infiltration of Goutallier grade 3 or higher was associated with worse post-operative outcomes and decreased active ER.

Many studies have evaluated the application of the two-incision technique popularised by Gerber.⁸ In the two-incision technique, a superior incision is made over the acromion. The deltoid is removed with a small piece of bone and reflected laterally. A humeral head devoid of posterosuperior tendon attachment is exposed, and

assessment of the remaining torn rotator cuff is performed. If any remaining rotator cuff can be incorporated into the transfer, it will be added. After preparation of the humeral head is performed, a second incision is made in the axilla. This incision starts in the mid-body of the latissimus dorsi and extends superior toward the axilla and then slants obliquely after the armpit. Careful dissection is necessary to find where the tendon inserts into the humerus. It is cut sharply off the humerus and a non-absorbable suture is placed in a Krakow fashion. The tendon is passed between the deltoid and the teres minor and is secured on the greater tuberosity as far anterior as possible.⁸ Gerber published his 10-year follow-up showing durability of the reconstruction, with 74% good to excellent results and maintenance of subject shoulder values and constant scores.³³ El-Azab et al confirmed this finding as well, demonstrating long-term failure rates of 10% and conversion to reverse shoulder arthroplasty (RSA) at 4%.¹⁹ Although the tenodesis effect may lose strength over time, it is possible to delay RSA after LDT.³⁴ Complications include stiffness, traumatic failure of the transfer, resolving nerve dysesthesia and deltoid re-attachment failure.³³

Techniques for LDT are evolving. Habermeyer³⁵ described a single-incision approach that uses a more posterior attachment of the transfer into the humeral head. Hertzberg et al¹¹ demonstrated that securing the transfer at the insertion of the infraspinatus improves rotational movement when compared with other locations. Instead of making an incision down the arm, the single-incision technique uses a V-shaped incision and direct visualisation of the posterior humeral head. One downside to this approach is that the anterior humeral head cannot readily be evaluated and thus small underlying subscapularis tears cannot be repaired. The published results were comparable with the Gerber two-incision technique regarding constant score, FF and ER. Recently, there have been advances in arthroscopically-assisted LDT.^{36,37,38,39} In a multi-centered study, Grimberg et al³⁷ demonstrated results equivalent to historical open two-incision approaches. They concluded that the one-incision arthroscopically-assisted approach provided better mechanical resistance to traction. Additionally, Gerhardt et al⁴⁰ did a five-year follow-up for the modified L'Episcopo⁴¹ single-incision technique which transferred both the latissimus dorsi and teres major. They demonstrated maintenance of constant scores at five years. However, there was persistent progression of cuff arthropathy. The LDTMT is popular when performed in combination with RSA for a positive ER lag sign.⁴²

Teres major transfers

Isolated teres major transfers (TMT) for irreparable posterosuperior RCTs are not well represented in the literature. It was designed as a transfer for infraspinatus deficiency.⁴³ This technique is performed using a two-incision technique similar to the previously described LDT

by Gerber. The teres major is isolated from the scapular origin from the latissimus dorsi. It is traced laterally to the humerus and removed with a strip of periosteum. Then it is passed under the deltoid and secured to the greater tuberosity. Interestingly, this transfer typically has an intact supraspinatus. If the supraspinatus is in jeopardy, it could be paired with a trapezius transfer. In 1998, Celli et al⁴³ were the first to publish results on this technique. Their case series of six patients demonstrated patient satisfaction with the operation. The same group published the long-term results of 20 patients with maintenance of improved post-operative constant scores.⁴⁴ A recent study was performed to evaluate the neurovascular responsibility to the muscle.⁴⁵ It was found that the lower subscapularis nerve innervates the teres major in over 85% of cases, with the difference supplied by the thoracodorsal nerve. The vascular supply inserts directly into the muscle within 2 cm of the nearby nerves, in the middle of the muscle belly. Often shorter in length than local nerves, the vascular structures tend to be the limiting factor when performing a TMT. In a biomechanical study, Steenbrink described improvement of post-operative activation when compared with LDT,⁴⁶ concluding that TMT was a superior anatomical transfer.

More recently, Henseler et al published short-term results of TMT transfers.⁴⁷ At two years, patients had improved FF, ER, visual analogue scale (VAS), and constant scores. Mansat et al⁴⁸ evaluated 12 TMTs with a mean of six years follow-up with similar results to Henseler. Similar to the LDT, negative prognostic factors included previous surgery and RCTs involving the subscapularis. Isolated infraspinatus involvement and a functional teres minor were positive prognostic factors. Further, Mansat et al described the following recommendations for TMT: the patient should be under the age of 55 years with a proper understanding of the condition and treatment, and an intact subscapularis and anterior supraspinatus cable.

LTT

LTT is a recently described modality for intervention of irreparable posterosuperior RCTs, and is a popular procedure for paralytic shoulders lacking ER.⁴⁹⁻⁵² Similar to TMT, LTT has not been well described in the literature. In this technique, a two-incision approach is performed as well. The first incision is made similar to the LDT, a sabre incision with deltoid take-down to expose the humeral head. The second incision is based 1 cm medial to the scapula. The lower trapezius attachment to the scapula is released and an Achilles allograft is used to augment for length. A subcutaneous tunnel is created from the medial incision to the lateral incision deep to the deltoid. A tendon wrap can be used to aid in gliding of the transfer. The transfer is secured to the footprint of the supraspinatus and upper border of the infraspinatus.

Most studies are anatomical or biomechanical. The lower trapezius tendon can be safely harvested without injury to the spinal accessory nerve or the muscle fibres of

the middle trapezius.⁵³ In a cadaveric study, Omid et al concluded that the LTT was superior to LDT for restoration of GH mechanics and joint reaction forces.⁵⁴ Hartzler et al⁵⁵ found that LTT had improved ER with the arm at the side compared with the LDT. In a poster presentation at the 2014 American Association of Orthopaedic Surgeons,⁵⁶ Galatz presented her series of 12 patients at a mean of 18 months follow-up. Significant improvement was noted in American Shoulder and Elbow Surgeons and VAS scores, FF, and ER. In total 75% of those examined with ultrasound demonstrated an intact transfer. The ER was noted to be better than historically noted for a LDT. In a recently accepted manuscript, Elhassan et al⁵⁷ followed 32 patients for a minimum of two years and found greater outcomes correlated with pre-operative FF > 60°. The most common complication of the transfer was a seroma owing to the necessary subcutaneous tunneling of the transfer.

Irreparable anterosuperior RCTs

Anterosuperior RCTs, more specifically subscapularis tears, occur far less commonly than the aforementioned posterosuperior RCTs.^{58,59} An MRI study of over 2000 RCTs demonstrated an incidence of 2% for subscapularis tears.⁶⁰ This number can be as high as 13% for isolated subscapularis tears and 37% of the time in combination with a supraspinatus tear.⁶¹ The subscapularis is essential for proper function of the shoulder as it is the primary internal rotator and anterior dynamic stabiliser.⁶² Patients with subscapularis RCTs typically present with anterior shoulder pain, internal rotation weakness, and dysfunction. Diminished strength can be assessed with multiple tests including the belly-press,⁶³ bear hug,⁶⁴ and lift-off.⁶⁵ While the repair of acute tears of the subscapularis muscle typically do well,⁶⁶⁻⁷⁰ the outcomes of chronic tears with fatty infiltration, Goutallier grade 3 or greater, are not as promising.^{58,71-73}

Excluding shoulder arthroplasty, several options for irreparable anterosuperior RCTs have been studied, including: pectoralis major transfer (PMT), pectoralis minor transfer (PMnT), latissimus transfer and teres major transfer.

PMT

The purpose of a PMT is to stand in for the subscapularis by exerting an internal rotation centering force.⁹ Wirth and Rockwood originally described the PMT in 1997.⁷² The PMT was performed anterior to the conjoined tendon and resulted in a high satisfaction rate at five years of follow-up. Resch et al⁷¹ adapted this technique to transfer only the superior two-thirds of the tendon under the conjoined tendon. Subsequently, Warner transferred the inferior sternal head attachment under the clavicular head but anterior to the conjoined tendon to avoid injury to the musculocutaneous nerve.⁷⁴

Biomechanically, subcoracoid placement of the transfer is superior.⁷⁵ However, there have been no comparative studies to date. Galatz et al⁷⁶ demonstrated a subcoracoid

transfer with over 75% satisfaction at almost 18 months with improved ASES and VAS scores. Elhassan et al evaluated patients treated with Warner's technique and found that those with a pre-operatively concentric humeral head had better outcomes.⁷⁷ Other recommendations include: age < 65 years (ideally < 40),⁷⁷ intact or reparable posterosuperior RCTs, and minimal GH arthritis.⁷⁸ It was noted that if the subcoracoid transfer is successful, ER loss up to 25° can be expected owing to a tenodesis effect.⁷¹

PMnT

Supratoracoid PMT can leave patients with a positive belly-press and lift-off test.⁷⁹ Investigators looked at other interventions including a PMnT.^{72,80} Wirth initially described this transfer in 1997 while reporting outcomes on both PMT and PMnT. Paladini used a subcoracoid PMnT with a small cortical piece of the coracoid for subscapularis tears of the superior two-thirds concomitant with irreparable supraspinatus tears. Two-year follow-up of 27 patients demonstrated significant improvement in constant score and FF. In total 78% returned to their activities of daily living. Interestingly, there was only an ER loss of 11° and a 22% presence of a belly-press test.

LDTMT

Cadaveric studies have been conducted in an effort to identify a more anatomical tendon transfer modality. Elhassan et al⁸¹ investigated the potential of LDT, TMT and LDTMT for anterior deficiencies. They concluded that a LDT or the TMT would be a viable safe option for transfer to the lesser tuberosity. *In vivo* studies must be conducted to determine the anatomical plausibility of these techniques.

Transfers for RTSA

Modern RTSA has provided surgeons with the ability to improve shoulder dysfunction in the setting of irreparable RCTs in the elderly.^{82,83,84} While the deltoid is the primary elevator of the arm, ER will be lacking if pre-operative evaluation demonstrated a positive Hornblower's or ER lag sign.^{82,85-87} The incidence is approximately 10%.⁸⁷ In an effort to restore ER, transfers using previously covered techniques have been studied, including LDT and LDTMT.

LDT

Gerber first described the two-incision LDT for RTSA in 2007.⁸⁸ Early results were promising; however, durability of the transfer was unknown. Puskas et al⁸⁹ evaluated 32 of Gerber's shoulders at intermediate-term follow-up and found excellent outcomes at five years with maintained ER. Surprisingly, the status of the teres minor had no effect on the outcome. It was concluded that patients with a pre-operative ER deficiency would benefit from a LDT in the setting of a RTSA.

LDTMT

In line with Gerber, Boileau published early results for his single-incision modified L'Episcopo technique where a LDTMT was performed.^{90,91} Boileau then prospectively evaluated 17 of his patients for at least one year after performing the aforementioned LDTMT.⁹² It was determined that the single-incision approach was easier, had fewer complications, and provided greater ER owing to addition of the TMT. Boughebi et al⁸⁷ re-affirmed these results with a minimum two-year follow-up investigation. Of note in both types of transfers (LDT or LDTMT), advanced age did not preclude the ability to perform either procedure.

Conclusion

Irreparable RCTs, Goutallier grade 3 or higher, can be debilitating. However, brilliant minds have developed techniques to improve quality of life and function. For posterosuperior RCTs, lacking the ability to raise and externally rotate the arm, the LDT remains the most time-tested transfer. LDT outcomes are satisfactory; however, LTT transfers are a promising new frontier and should be considered. Isolated subscapularis deficiencies seem to be well treated with PMT. In the presence of an irreparable supraspinatus and anterior supraspinatus one can consider the PMnT. When presented with rotator cuff arthropathy and an ER lag, the single-incision LDTMT appears to be the 'go to' transfer, as it improves ER with fewer complications compared with the isolated LDT.

AUTHOR INFORMATION

First State Orthopaedics, Newark, USA

Correspondence should be sent to Jeremie M. Axe, First State Orthopaedics, 4745 Stanton-Ogletown Road, Suite 225 Newark, DE 19713, USA.

E-mail: jeremieaxe@gmail.com

CONFLICT OF INTEREST

None declared.

FUNDING

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

LICENCE

© 2016 The author(s)

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) licence (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed.

REFERENCES

1. Bartl C, Kouloumentas P, Holzapfel K, et al. Long-term outcome and structural integrity following open repair of massive rotator cuff tears. *Int J Shoulder Surg* 2012;6:1-8. <https://creativecommons.org/licenses/by-nc/4.0/>

2. **Jo CH, Shin JS, Lee YG, et al.** Platelet-rich plasma for arthroscopic repair of large to massive rotator cuff tears: a randomized, single-blind, parallel-group trial. *Am J Sports Med* 2013;41:2240-2248.
3. **Kim SJ, Kim SH, Lee SK, Seo JW, Chun YM.** Arthroscopic repair of massive contracted rotator cuff tears: aggressive release with anterior and posterior interval slides do not improve cuff healing and integrity. *J Bone Joint Surg [Am]* 2013;95-A:1482-1488.
4. **Shamsudin A, Lam PH, Peters K, et al.** Revision versus primary arthroscopic rotator cuff repair: a 2-year analysis of outcomes in 360 patients. *Am J Sports Med* 2015;43:557-564.
5. **Iannotti JP, Deutsch A, Green A, et al.** Time to failure after rotator cuff repair: a prospective imaging study. *J Bone Joint Surg [Am]* 2013;95-A:965-971.
6. **Koh KH, Laddha MS, Lim TK, Park JH, Yoo JC.** Serial structural and functional assessments of rotator cuff repairs: do they differ at 6 and 19 months postoperatively? *J Shoulder Elbow Surg* 2012;21:859-866.
7. **Parnes N, DeFranco M, Wells JH, Higgins LD, Warner JJ.** Complications after arthroscopic revision rotator cuff repair. *Arthroscopy* 2013;29:1479-1486.
8. **Gerber C, Vinh TS, Hertel R, Hess CW.** Latissimus dorsi transfer for the treatment of massive tears of the rotator cuff. A preliminary report. *Clin Orthop Relat Res* 1988;232:51-61.
9. **Omid R, Lee B.** Tendon transfers for irreparable rotator cuff tears. *A Am Acad. Orthop Surg* 2013;21:492-501.
10. **Schmidt CC, Jarrett CD, Brown BT.** Management of rotator cuff tears. *J Hand Surg Am* 2015;40:399-408.
11. **Herzberg G, Urien JP, Dimnet J.** Potential excursion and relative tension of muscles in the shoulder girdle: relevance to tendon transfers. *J Shoulder Elbow Surg* 1999;8:430-437.
12. **Namdari S, Voleti P, Baldwin K, Glaser D, Huffman GR.** Latissimus dorsi tendon transfer for irreparable rotator cuff tears: a systematic review. *J Bone Joint Surg [Am]* 2012;94-A:891-898.
13. **Warner JJ, Parsons IM IV.** Latissimus dorsi tendon transfer: a comparative analysis of primary and salvage reconstruction of massive, irreparable rotator cuff tears. *J Shoulder Elbow Surg* 2001;10:514-521.
14. **Gerber C, Maquieira G, Espinosa N.** Latissimus dorsi transfer for the treatment of irreparable rotator cuff tears. *J Bone Joint Surg [Am]* 2006;88-A:113-120.
15. **Irlenbusch U, Bracht M, Gansen HK, Lorenz U, Thiel J.** Latissimus dorsi transfer for irreparable rotator cuff tears: a longitudinal study. *J Shoulder Elbow Surg* 2008;17:527-534.
16. **Moursy M, Forstner R, Koller H, Resch H, Tauber M.** Latissimus dorsi tendon transfer for irreparable rotator cuff tears: a modified technique to improve tendon transfer integrity. *J Bone Joint Surg [Am]* 2009;91-A:1924-1931.
17. **Nové-Josserand L, Costa P, Liotard JP, et al.** Results of latissimus dorsi tendon transfer for irreparable cuff tears. *Orthop Traumatol Surg Res* 2009;95:108-113.
18. **Gerber C.** Latissimus dorsi transfer for the treatment of irreparable tears of the rotator cuff. *Clin Orthop Relat Res* 1992;275:152-160.
19. **El-Azab HM, Rott O, Irlenbusch U.** Long-term follow-up after latissimus dorsi transfer for irreparable posterosuperior rotator cuff tears. *J Bone Joint Surg [Am]* 2015;18:97-A:462-469.
20. **Glanzmann MC, Goldhahn J, Flury M, Schwyzer HK, Simmen BR.** Deltoid flap reconstruction for massive rotator cuff tears: mid- and long-term functional and structural results. *J Shoulder Elbow Surg* 2010;19:439-445.
21. **Werner CM, Zingg PO, Lie D, Jacob HA, Gerber C.** The biomechanical role of the subscapularis in latissimus dorsi transfer for the treatment of irreparable rotator cuff tears. *J Shoulder Elbow Surg* 2006;15:736-742.
22. **McFarland EG.** *Examination of the shoulder: the complete guide.* New York: Thieme, 2006.
23. **Hertel R, Ballmer FT, Lombert SM, Gerber C.** Lag signs in the diagnosis of rotator cuff rupture. *J Shoulder Elbow Surg* 1996;5:307-313.
24. **Buchmann S, Plath JE, Imhoff AB.** Latissimus dorsi transfer for the treatment of irreparable rotator tears: indication, surgical technique, and modifications. *Oper Orthop Traumatol* 2012;24:502-512. (In German).
25. **Iannotti JP, Hennigan S, Herzog R, et al.** Latissimus dorsi tendon transfer for irreparable posterosuperior rotator cuff tears. Factors affecting outcome. *J Bone Joint Surg [Am]* 2006;88-A:342-348.
26. **Hamada K, Yamanaka K, Uchiyama Y, Mikasa T, Mikasa M.** A radiographic classification of massive rotator cuff tear arthritis. *Clin Orthop Relat Res* 2011;469:2452-2460.
27. **Patte D.** Classification of rotator cuff lesions. *Clin Orthop Relat Res* 1990;254:81-86.
28. **Goutallier D, Postel JM, Bernageau J, Lavau L, Voisin MC.** Fatty infiltration of disrupted rotator cuff muscles. *Rev Rhum Engl Ed* 1995;62:415-422.
29. **Fuchs B, Weishaupt D, Zanetti M, Hodler J, Gerber C.** Fatty degeneration of the muscles of the rotator cuff: assessment by computed tomography versus magnetic resonance imaging. *J Shoulder Elbow Surg* 1999;8:599-605.
30. **Thomazeau H, Rolland Y, Lucas C, Duval JM, Langlais F.** Atrophy of the supraspinatus belly. Assessment by MRI in 55 patients with rotator cuff pathology. *Acta Orthop Scand* 1996;67:264-268.
31. **Fox J, Romeo AA.** Arthroscopic subscapularis repair. In: *Annual meeting of the American Academy of Orthopaedic Surgeons*, 2003. New Orleans, LA.
32. **Costouros JG, Espinosa N, Schmid MR, Gerber C.** Teres minor integrity predicts outcome of latissimus dorsi tendon transfer for irreparable rotator cuff tears. *J Shoulder Elbow Surg* 2007;16:727-734.
33. **Gerber C, Rahm SA, Catanzaro S, Farshad M, Moor BK.** Latissimus dorsi tendon transfer for treatment of irreparable posterosuperior rotator cuff tears: long-term results at a minimum follow-up of ten years. *J Bone Joint Surg [Am]* 2013;95-A:1920-1926.
34. **Erşen A, Ozben H, Demirhan M, Atalar AC, Kapıcıoğlu M.** Time-dependent changes after latissimus dorsi transfer: tenodesis or tendon transfer? *Clin Orthop Relat Res* 2014;472:3880-3888.
35. **Habermeyer P, Magosch P, Rudolph T, Lichtenberg S, Liem D.** Transfer of the tendon of latissimus dorsi for the treatment of massive tears of the rotator cuff: a new single-incision technique. *J Bone Joint Surg [Br]* 2006;88:208-212.
36. **Grimberg J, Kany J, Valenti P, Amaravathi R, Ramalingam AT.** Arthroscopic-assisted latissimus dorsi tendon transfer for irreparable posterosuperior cuff tears. *Arthroscopy* 2015;31:599-607.e1.
37. **Grimberg J, Kany J.** Latissimus dorsi tendon transfer for irreparable postero-superior cuff tears: current concepts, indications, and recent advances. *Curr Rev Musculoskelet Med* 2014;7:22-32.
38. **Villacis D, Merriman J, Wong K, Rick Hatch GF III.** Latissimus dorsi tendon transfer for irreparable rotator cuff tears: a modified technique using arthroscopy. *Arthrosc Tech* 2013;2:e27-e30.
39. **Castricini R, Longo UG, De Benedetto M, et al.** Arthroscopic-assisted latissimus dorsi transfer for the management of irreparable rotator cuff tears: Short-term results. *J Bone Joint Surg [Am]* 2014;96-A:e119.
40. **Gerhardt C, Lehmann L, Lichtenberg S, Magosch P, Habermeyer P.** Modified L'Episcopo tendon transfers for irreparable rotator cuff tears: 5-year follow-up. *Clin Orthop Relat Res* 2010;468:1572-1577

41. **Habermeyer P, Magosch P, Lichtenberg S.** The modified L'Episcopo procedure to reconstruct massive rotator cuff tears — A prospective study. In: *Annual meeting of the American Academy of Orthopaedic Surgeons*, 2002. Dallas, TX.
42. **Boileau P, Chuinard C, Roussanne Y, Neyton L, Trojani C.** Modified latissimus dorsi and teres major transfer through a single delto-pectoral approach for external rotation deficit of the shoulder: as an isolated procedure or with a reverse arthroplasty. *J Shoulder Elbow Surg* 2007;16:671-682.
43. **Celli L, Rovesta C, Marongiu MC, Manzieri S.** Transplantation of teres major muscle for infraspinatus muscle in irreparable rotator cuff tears. *J Shoulder Elbow Surg* 1998;7:485-490.
44. **Celli A, Marongiu MC, Rovesta C, Celli L.** Transplant of the teres major in the treatment of irreparable injuries of the rotator cuff (long-term analysis of results). *Chir Organi Mov* 2005;90:121-132. (In English, Italian).
45. **Dancker M, Lambert S, Brenner E.** The neurovascular anatomy of the teres major muscle. *J Shoulder Elbow Surg* 2015;24:e57-e67.
46. **Steenbrink F, Nelissen RG, Meskers CG, et al.** Teres major muscle activation relates to clinical outcome in tendon transfer surgery. *Clin Biomech (Bristol, Avon)* 2010;25:187-193.
47. **Henseler JF, Nagels J, van der Zwaal P, Nelissen RG.** Teres major tendon transfer for patients with massive irreparable posterosuperior rotator cuff tears: short-term clinical results. *Bone Joint J* 2013;95-B:523-529.
48. **Mansat P, Dotziz A, Bellumore Y, Mansat M.** Teres major flap: surgical anatomy, technique of harvesting, methods of fixation, postoperative management. In: Valenti P, ed. *Tendon transfer for irreparable rotator cuff tear*. Paris: Springer-Verlag France, 2011:49-64.
49. **Elhassan B, Bishop A, Shin A, Spinner R.** Shoulder tendon transfer options for adult patients with brachial plexus injury (review). *J Hand Surg Am* 2010;35-A:1211-1219.
50. **Elhassan B, Bishop A, Shin A.** Trapezius transfer to restore external rotation in a patient with a brachial plexus injury. A case report. *J Bone Joint Surg [Am]* 2009;91-A:939-944.
51. **Elhassan B, Bishop AT, Hartzler RU, Shin AY, Spinner RJ.** Tendon transfer options about the shoulder in patients with brachial plexus injury. *J Bone Joint Surg [Am]* 2012;94-A:1391-1398.
52. **Elhassan B.** Technique of tendon transfers about the shoulder in patients with brachial plexus injury. *JBJS Essential Surgical Techniques* 2012;2:e19.1-12.
53. **Omid R, Cavallero MJ, Granholm D, Villacis DC, Yi AM.** Surgical anatomy of the lower trapezius tendon transfer. *J Shoulder Elbow Surg* 2015;24:1353-1358.
54. **Omid R, Heckmann N, Wang L, et al.** Biomechanical comparison between the trapezius transfer and latissimus transfer for irreparable posterosuperior rotator cuff tears. *J Shoulder Elbow Surg* 2015;24:1635-1643.
55. **Hartzler RU, Barlow JD, An KN, Elhassan BT.** Biomechanical effectiveness of different types of tendon transfers to the shoulder for external rotation. *J Shoulder Elbow Surg* 2012;21:1370-1376.
56. **Donegan RP, Jobin CM, Chamberlain AM, et al.** Lower trapezius tendon transfer for irreparable posterior-superior rotator cuff tears. In: *Annual meeting of the American Academy of Orthopaedic Surgeons*, 2014; New Orleans, LA.
57. **Elhassan B, Wagner E, Werthel JD.** Outcome of lower trapezius transfer to reconstruct massive irreparable posterior-superior rotator cuff tear. *J Shoulder Elbow Surg* 2015. (Epub ahead of print).
58. **Lyons RP, Green A.** Subscapularis tendon tears. *J Am Acad Orthop Surg* 2005;13:353-363.
59. **Sakurai G, Ozaki J, Tomita Y, Kondo T, Tamai S.** Incomplete tears of the subscapularis tendon associated with tears of the supraspinatus tendon: cadaveric and clinical studies. *J Shoulder Elbow Surg* 1998;7:510-515.
60. **Li XX, Schweitzer ME, Bifano JA, et al.** MR evaluation of subscapularis tears. *J Comput Assist Tomogr* 1999;23:713-717.
61. **Lanz U, Fullick R, Bongiorno V, et al.** Arthroscopic repair of large subscapularis tendon tears: 2- to 4-year clinical and radiographic outcomes. *Arthroscopy* 2013;29:1471-1478.
62. **Perry J.** Anatomy and biomechanics of the shoulder in throwing, swimming, gymnastics, and tennis. *Clin Sports Med* 1983;2:247-270.
63. **Burkhart SS, Tehrany AM.** Arthroscopic subscapularis tendon repair: technique and preliminary results. *Arthroscopy* 2002;18:454-463.
64. **Barth JR, Burkhart SS, De Beer JF.** The bear-hug test: a new and sensitive test for diagnosing a subscapularis tear. *Arthroscopy* 2006;22:1076-1084.
65. **Gerber C, Krushell RJ.** Isolated rupture of the tendon of the subscapularis muscle. Clinical features in 16 cases. *J Bone Joint Surg [Br]* 1991;73-B:389-394.
66. **Adams CR, Schoolfield JD, Burkhart SS.** The results of arthroscopic subscapularis tendon repairs. *Arthroscopy* 2008;24:1381-1389.
67. **Bartl C, Salzmann GM, Seppel G, et al.** Subscapularis function and structural integrity after arthroscopic repair of isolated subscapularis tears. *Am J Sports Med* 2011;39:1255-1262.
68. **Bartl C, Scheibel M, Magosch P, Lichtenberg S, Habermeyer P.** Open repair of isolated traumatic subscapularis tendon tears. *Am J Sports Med* 2011;39:490-496.
69. **Denard PJ, Jiwani AZ, Lädemann A, Burkhart SS.** Long-term outcome of a consecutive series of subscapularis tendon tears repaired arthroscopically. *Arthroscopy* 2012;28:1587-1591.
70. **Ide J, Tokiyoshi A, Hirose J, Mizuta H.** Arthroscopic repair of traumatic combined rotator cuff tears involving the subscapularis tendon. *J Bone Joint Surg [Am]* 2007;89-A:2378-2388.
71. **Resch H, Povacz P, Ritter E, Matschi W.** Transfer of the pectoralis major muscle for the treatment of irreparable rupture of the subscapularis tendon. *J Bone Joint Surg [Am]* 2000;82-A:372-382.
72. **Wirth MA, Rockwood CA Jr.** Operative treatment of irreparable rupture of the subscapularis. *J Bone Joint Surg [Am]* 1997;79-A:722-731.
73. **Warner JP, Gerber C.** Treatment of massive rotator cuff tears: posterior-superior and anterior-superior. In: Iannotti JP, ed. *The rotator cuff: current concepts and complex problems*. Rosemont, Illinois: American Academy of Orthopedic Surgeons, 1998:59-94.
74. **Warner JJ.** Management of massive irreparable rotator cuff tears: the role of tendon transfer. *Instr Course Lect* 2001;50:63-71.
75. **Konrad GG, Sudkamp NP, Kreuz PC, et al.** Pectoralis major tendon transfers above or underneath the conjoint tendon in subscapularis-deficient shoulders. An in vitro biomechanical analysis. *J Bone Joint Surg [Am]* 2007;89-A:2477-2484.
76. **Klepps S, Galatz LM, Yamaguchi K.** Subcoracoid pectoralis major transfer: A salvage procedure for irreparable subscapularis deficiency. *Tech Shoulder Elbow Surg* 2001;2:85-91.
77. **Elhassan B, Ozbaydar M, Massimini D, et al.** Transfer of pectoralis major for the treatment of irreparable tears of subscapularis: does it work? *J Bone Joint Surg [Br]* 2008;90-B:1059-1065.
78. **Nelson GN, Namdari S, Galatz L, Keener JD.** Pectoralis major tendon transfer for irreparable subscapularis tears. *J Shoulder Elbow Surg* 2014;23:909-918.

- 79. Jost B, Puskas GJ, Lustenberger A, Gerber C.** Outcome of pectoralis major transfer for the treatment of irreparable subscapularis tears. *J Bone Joint Surg[Am]* 2003; 85-A:1944-1951.
- 80. Paladini P, Campi F, Merolla G, Pellegrini A, Porcellini G.** Pectoralis minor tendon transfer for irreparable anterosuperior cuff tears. *J Shoulder Elbow Surg* 2013;22:e1-e5.
- 81. Elhassan B, Christensen TJ, Wagner ER.** Feasibility of latissimus and teres major transfer to reconstruct irreparable subscapularis tendon tear: an anatomic study. *J Shoulder Elbow Surg* 2014;23:492-499.
- 82. Boileau P, Watkinson DJ, Hatzidakis AM, Balg F.** Grammont reverse prosthesis: design, rationale, and biomechanics. *J Shoulder Elbow Surg* 2005;14(1 suppl S):147S-161S.
- 83. Boulahia A, Edwards TB, Walch G, Baratta RV.** Early results of a reverse design prosthesis in the treatment of arthritis of the shoulder in elderly patients with a large rotator cuff tear. *Orthopedics* 2002;25:129-133.
- 84. Jacobs R, Debeer P, De Smet L.** Treatment of rotator cuff arthropathy with a reversed Delta shoulder prosthesis. *Acta Orthop Belg* 2001;67:344-347.
- 85. Walch G, Boulahia A, Calderone S, Robinson AH.** The 'dropping' and 'hornblower's' signs in evaluation of rotator-cuff tears. *J Bone Joint Surg[Br]* 1998;80-B:624-628.
- 86. Simovitch RW, Helmy N, Zumstein MA, Gerber C.** Impact of fatty infiltration of the teres minor muscle on the outcome of reverse total shoulder arthroplasty. *J Bone Joint Surg [Am]* 2007;89-A:934-939.
- 87. Boughebri O, Kilinc A, Valenti P.** Reverse shoulder arthroplasty combined with a latissimus dorsi and teres major transfer for a deficit of both active elevation and external rotation. Results of 15 cases with a minimum of 2-year follow-up. *Orthop Traumatol Surg Res* 2013;99:131-137.
- 88. Gerber C, Pennington SD, Lingenfelter EJ, Sukthankar A.** Reverse Delta-III total shoulder replacement combined with latissimus dorsi transfer. A preliminary report. *J Bone Joint Surg[Am]* 2007;89-A:940-947.
- 89. Puskas GJ, Catanzaro S, Gerber C.** Clinical outcome of reverse total shoulder arthroplasty combined with latissimus dorsi transfer for the treatment of chronic combined pseudoparesis of elevation and external rotation of the shoulder. *J Shoulder Elbow Surg* 2014;23:49-57.
- 90. Boileau P, Chuinard C, Roussanne Y, et al.** Reverse shoulder arthroplasty combined with a modified latissimus dorsi and teres major tendon transfer for shoulder pseudoparesis associated with dropping arm. *Clin Orthop Relat Res* 2008;466:584-593.
- 91. Boileau P, Chuinard C, Roussanne Y, Neyton L, Trojani C.** Modified latissimus dorsi and teres major transfer through a single delto-pectoral approach for external rotation deficit of the shoulder: as an isolated procedure or with a reverse arthroplasty. *J Shoulder Elbow Surg* 2007;16:671-682.
- 92. Boileau P, Rumian AP, Zumstein MA.** Reversed shoulder arthroplasty with modified L'Episcopo for combined loss of active elevation and external rotation. *J Shoulder Elbow Surg* 2010;19(2 suppl):20-30.